

Event horizons and apparent horizons in spherically symmetric geometries.

Edward Malec

Physics Department, UCC, Cork, Ireland and

Institute of Physics, UJ 30-059 Cracow, Reymonta 4, Poland

Spherical configurations that are very massive must be surrounded by apparent horizons. These in turn, when placed outside a collapsing body, have a fixed area and must propagate outward with a velocity equal to the velocity of radially outgoing photons. That proves, within the framework of the (1+3) formalism and without resorting to the Birkhoff theorem, that apparent horizons coincide with event horizons in electrovacuum.

1. Introduction.

The Cosmic Censorship Hypothesis (CCH) [?] is certainly the most compelling open problem of classical general relativity. From the dynamical point of view any successful attempt to prove a weak version of CCH in a space-time generated by an isolated self-gravitating object must consist of the following points:

i) state smooth initial data (if this is done then space-time can be unambiguously splitted into time- and space- directions, initially at least, once the local Cauchy problem is solved);

ii) prove that if a singularity emerges then it must be hidden inside an event horizon, so as not to influence the asymptotically flat open end;

iii) prove the global Cauchy problem in the asymptotically flat region outside the event horizon (this implies that space-time splitting is possible globally in the region and we are guaranteed the existence of the line element inside it).

From this perspective the proof of CCH seems to be technically unattainable (at least nowadays) in the general case of a nonspherical collapse, since the global Cauchy problem is almost intractable at present [?]. Even in the spherically symmetric collapse only partial results are known [?].

It is natural to assume the solvability of the global Cauchy problem, that is to assume the global existence of a space-time metric, in order to test the remaining steps of the above programme. In this case Israel has proven the confining property of apparent horizons [?] in spherically symmetric geometries [?]. Israel's result opens a way to prove CCH for those singularities that must be hidden inside apparent horizons. The formation of apparent horizons, in turn, in spherically symmetric space-times has been completely solved in the initial value formulation [?], [?], thus accomplishing the proof of steps i) and ii) for a version of CCH in spherically symmetric geometries.

The intention of this paper is to provide another proof of the Israel's result in the framework of a dynamical (1+3) description of a spherical collapse. I prove a confin-

ing property of apparent horizons that are placed in electrovacuum and complement this with a proof that a relevant solution of the global Cauchy problem exists.

The confining property of apparent horizons in electrovacuum is not a surprise. It is well known thanks to the Birkhoff theorem [?] that once the areal radius $\mathbf{R}$ of a charged collapsing body becomes equal to $m + \sqrt{m^2 - q^2}$, where m is the asymptotic (Einstein - Freund - Arnowitt - Deser - Misner) mass and q is the total charge, then the body hides within an event horizon that coincides with a sphere of the areal radius $\mathbf{R}$ which in turn is the locus of an apparent horizon. This conclusion appears correct despite the fact that the Birkhoff transformation does not exist in the situation of interest when a geometry contains apparent horizons (see a discussion in the Appendix of [?] in which a generalized version of the Birkhoff theorem is described).

In the first two sections I will assume the existence of a global Cauchy solution, that is the existence of an asymptotically flat space-time with the spherically symmetric metric line element

$$ds^2 = -\alpha^2(r, t)dt^2 + a(r, t)dr^2 + b(r, t)r^2d\Omega^2. \quad (1)$$

r is a coordinate radius and $br^2d\Omega^2$ is a standard 2-sphere metric element. We assume the maximal gauge condition in which components of the extrinsic curvature K_{ij} of the hypersurface Σ_t (defined as a set of points having a fixed coordinate time $t = \text{const}$) satisfy the following equations

$$K_r^r = \frac{\partial_t a}{2a |\alpha|} = -2K_\phi^\phi = -2K_\theta^\theta = -\frac{\partial_t b}{b |\alpha|}. \quad (2)$$

We use the standard convention [?] in which quantities supplied with Greek type indices refer to the four-geometry while quantities with Latin labels refer to the geometry of the hypersurface Σ_t . The Einstein summation convention is applied in some places, with the exception concerning the label r , whose repetition is assumed never to mean summation. Below ∇_i denotes the covariant derivative on Σ_t . The Einstein equations read

$$-\frac{2}{abr^2}\partial_r(r^2\partial_rb) + \frac{(\partial_rb)^2}{2ab^2} + \frac{\partial_rb\partial_ra}{a^2b} + \frac{2\partial_ra}{a^2r} - \frac{2\partial_rb}{abr} + \frac{2}{r^2}\left(\frac{1}{b} - \frac{1}{a}\right) = K_{ij}K^{ij} + 16\pi\rho. \quad (3)$$

$$\nabla_i K^{ir} = -8\pi j^r. \quad (4)$$

$$\nabla_i \partial^i \alpha = K_{ij}K^{ij}\alpha + 4\pi[-T_0^0 + T_i^i]\alpha \quad (5)$$

$$\partial_t K_r^r = -\alpha R_r^{(3)r} - 8\pi\alpha[-T_r^r + \frac{T_\mu^\mu}{2}] + \nabla_r \partial^r \alpha \quad (6)$$

Above T_μ^ν is the energy momentum tensor of matter generating the gravitational field, $\rho = -T_0^0$ and $j_i = -T_i^0\alpha$. $R_r^{(3)r}$ denotes a radial component of the 3-dimensional Ricci tensor which might be expressed (using the hamiltonian constraint) as follows

$$R_r^{(3)r} = -8\pi T_0^0 + \frac{K_{ij}K^{ij}}{2} + \frac{(\partial_rb)^2}{4ab^2} + \frac{\partial_rb}{abr} - \frac{1}{br^2} + \frac{1}{ar^2}. \quad (7)$$

The equations (??, ??) are dynamical ones while (??, ??) are the hamiltonian and momentum constraints [?], respectively. As initial data one may take for instance $a(r, 0) = b(r, 0)$ at a time $t = 0$; given the matter distribution one obtains from (??, ??) the three-geometry of Σ and the extrinsic curvature K_r^r and from (??) the lapse α . Equations (??, ??) determine the rate of change of the three-geometry and of K_r^r . Of the two functions a, b only one is independent. In the maximal slicing $\partial_t(b^2a) = 0$; taking into account the above initial condition one gets

$$b^2(r, t) = \frac{b^3(r, 0)}{a(r, t)}. \quad (8)$$

The order of the rest of this paper is following. In Section 2 it is proved that the apparent horizon is null-like in electrovacuum, has a constant area and coincides with the event horizon. Section 3 contains a proof of a version of the global Cauchy problem. In the last Section I comment on the significance of the results and their (possible) generalization.

2. Main calculations.

Let the boundary of a collapsing body be a sphere of a coordinate radius r_0 . Let us define $c = \frac{-q^2+m^2}{4}$. First, let us notice that there exists a solution (Appendix 1) that is manifestly static outside a collapsing body [?]

$$a = b = \left(1 + \frac{m}{r} + \frac{c}{r^2}\right)^2, \quad (9)$$

$$|\alpha| = \left| \frac{r - \sqrt{c}}{r + \sqrt{c}} \right|^{\frac{m}{2\sqrt{c}}} \left| r^2 - c \right|^{\frac{q^2}{2(r^2+mr+c)}} e^{\frac{q^2}{2} \int_r^\infty \frac{\ln(|s^2-c^2|)(2s+m)}{(s^2+ms+c)^2} ds}. \quad (10)$$

α vanishes at $r = \sqrt{c}$ quicker than $r - \sqrt{c}$, since $\frac{m}{2\sqrt{c}} \geq 1$. From this one readily infers that at a surface S placed at a coordinate radius $r = \sqrt{c}$ (that is, at an areal radius $\mathbf{R} = m + \sqrt{m^2 - q^2}$) exists an event horizon; no signal can traverse through S in a finite coordinate time t . In the case of vanishing total charge q the corresponding line element coincides with the Schwarzschild line element in isotropic coordinates [?]. The solution (??, ??) will not be considered in the rest of this paper.

Let us assume that a part of a spherically symmetric spacetime generated by a collapsing body can be foliated by maximal slices Σ_t , $0 \leq t \leq t_0$ that are asymptotically flat. Assume also that there exists a smooth continuation of the above band of slices, hypersurfaces Σ_t^{out} , that are maximal outside a region of compact support and that cover a region with the outermost apparent horizon (if it exists). The coordinate time t is a parameter that labels maximal slices but it coincides with a proper time of an external observer that is localized very far from a collapsing body.

It was proven elsewhere [?] that when the amount of matter minus a total radial momentum exceeds a multiple (1 or 7/6) of the proper radius, then apparent horizons must form. Let us assume that there exists an apparent horizon outside a (neutral or charged) collapsing body of a compact support. (We do not exclude electrovacuum, i. e., there might exist long-ranged potentials outside a body, with an electrostatic Coulomb-like energy density.)

Under these conditions, one proves that the Penrose [?], [?] inequality (which actually becomes an equality) holds true, that at the surface of an apparent horizon

$$m = \sqrt{\frac{S}{16\pi}} + q^2 \sqrt{\frac{\pi}{S}}. \quad (11)$$

It will be convenient to prove (??) in an isotropic system of coordinates in which $a = b = \phi^4$. It is easy to prove that this form of a metric can be achieved just by performing a suitable change of a radial coordinate on a fixed Cauchy slice. Moreover, the final result - equation (??) - is already expressed in a coordinate independent way.

The proof goes as follows. In electro-vacuum the hamiltonian constraint reads [?]

$$\hat{\Delta}\phi = -\frac{1}{4}\hat{E}_i\hat{E}^i\phi^{-3} - \frac{K_{ij}K^{ij}\phi^5}{8}. \quad (12)$$

Here the hatted quantities refer to the flat background metric and $\hat{E}^r = \frac{q}{r^2}, E^\theta = E^\phi = 0$. From the momentum constraints one gets [?]

$$K_{ij} = (n_i n_j - \frac{g_{ij}}{3}) \frac{C}{\phi^6 R^3}, \quad (13)$$

where n_j is a normal vector in the physical non-hatted metric. C depends on time but it is constant on a part of a fixed Cauchy slice that is exterior to the collapsing body.

Equation (??) has a conserved (r -independent, $\partial_r E = 0$) quantity

$$E = \frac{r}{8}(2r\partial_r\phi + \phi)^2 - \frac{r\phi^2}{8} - \frac{q^2}{8r\phi^2} - \frac{C^2}{72r^3\phi^6}. \quad (14)$$

Assuming asymptotic flatness one finds that $E = -\frac{m}{4}$, where m is the asymptotic mass. Notice also that $r\phi^2$ is equal to an areal radius, $r\phi^2 = \sqrt{\frac{S}{4\pi}}$. After some rearrangements one might write (??) as follows

$$m - [\sqrt{\frac{S}{16\pi}} + q^2 \sqrt{\frac{\pi}{S}}] = -\frac{r}{2}[2r\partial_r\phi + \phi - \frac{C}{3r^2\phi^3}][2r\partial_r\phi + \phi + \frac{C}{3r^2\phi^3}] = -\frac{\mathbf{R}^3}{8}\theta(S)\theta'(S). \quad (15)$$

Here $\theta(S)$ is the divergence of outgoing light rays, $\theta(S) = \frac{1}{\alpha} \frac{d}{dt} \ln S$ (the derivative in the direction of outgoing photons) and $\theta'(S) = -\frac{1}{\alpha} \frac{d}{dt} \ln S$ (the derivative in the

direction of incoming photons)⁷ is the convergence of incoming light rays. If S is an apparent horizon, then $\theta(S)$ vanishes which proves (??).

Equation (??) actually holds on all maximal slices or on their time developments that are maximal in an asymptotically flat region (up to the apparent horizons) as far as the apparent horizons remains outside the collapsing body. That means that the areal radius

$$\mathbf{R} = r\sqrt{b} \quad (16)$$

(we are coming back to the original metric notation (??)) of the apparent horizon must be conserved in time, since the mass m is conserved in time in asymptotically flat systems. That is, the full time derivative of $\mathbf{R}$ must vanish, which leads to the following equality

$$V\sqrt{b}\frac{2b + r\partial_r b}{2b} + r\frac{\partial_t b}{2\sqrt{b}} = 0. \quad (17)$$

Here $V = dr/dt$ is the coordinate velocity expansion of the apparent horizon. Now, the condition for the apparent horizon reads

$$2b + r\partial_r b = \frac{brK_{rr}}{\sqrt{a}}; \quad (18)$$

inserting this into (??) and using (??) we obtain

$$\frac{\sqrt{b}K_{rr}r}{1\sqrt{a}}[V - \frac{|\alpha|}{\sqrt{a}}] = 0. \quad (19)$$

We conclude that the apparent horizon expands with a radial velocity

$$V = \frac{|\alpha|}{\sqrt{a}}. \quad (20)$$

But, from equation (1) we know that $\frac{|\alpha|}{\sqrt{a}}$ is equal to the velocity of radially outgoing photons. Therefore, no material object can escape from within the apparent horizon; it just coincides with the event horizon. We can say more. Actually, the equation (??) might be interpreted in the following way: if an areal radius of a sphere S satisfies (??), then S is an apparent horizon. By continuity, all Cauchy slices must

contain surfaces satisfying (??), and the surfaces must remain in vacuum, since they move with the velocity of light. That means that, as long as maximal slices exist (or a time development of an initially maximal slice that is partially maximal later on, that is it remains maximal in an open end containing the apparent horizon) and under conditions stated previously, the apparent horizon must exist forever and it coincides with an event horizon. In this way we have proven in the framework of (1+3) formalism, without resorting to the Birkhoff theorem, that there exists an event horizon that intersects each Cauchy slice along a sphere of the areal radius $\mathbf{R} = m + \sqrt{m^2 - q^2}$.

Let us remark that one can prove in a similar way, that when a locus of points with $\theta' = 0$ (i. e., the past apparent horizon) is placed in electro-vacuum, then it moves to the centre with the velocity of light. This set of points is impenetrable from outside and its area is constant, hence it constitutes a boundary of a white hole [?]. The solution (??, ??) is simultaneously a white hole and a black hole since $\theta = \theta' = 0$ at $r = \sqrt{c}$. The velocity of the boundary $r = \sqrt{c}$ is equal to zero.

There exists yet another possibility to prove that if the evolution of a collapsing matter is smooth outside a region of compact support (the latter can contain singularities), then if an apparent horizon exists at a time t then it must exist forever. This is obvious, if we notice that the development of the divergence θ in the direction of outgoing photons is given by the Raychaudhury equation [?], which in electrovacuum reads in (1+3) splitting [?]

$$\frac{d}{dt}\theta = \frac{\theta}{\sqrt{a}}(\partial_r\alpha + \alpha\sqrt{a}K_r^r) - \frac{\alpha\theta^2}{2}. \quad (21)$$

From (??) one can deduce that a surface with vanishing divergence θ propagates to the future.

Equation (??) provides a well known necessary and (simultaneously) sufficient condition for the formation of event horizons formulated in terms of asymptotic quantities m, q and an area S . One can obtain also criteria in terms of quasilocal

quantities, by combining results of this paper with some of theorems of [?]. One can, for instance, formulate the following statement:

Theorem 1. If on an initial spherically symmetric Cauchy hypersurface with momentarily static initial data

$$M > L, \tag{22}$$

where M is a total mass [?] and L is a proper radius of a collapsing body, then the whole space-time history of the body must contain an event horizon that surrounds the body.

Thus, if the energy content inside a ball of a fixed radius becomes large, then it hides under an event horizon. That proves a version of the Cosmic Censorship Hypothesis [?] (CCH) in which *singularities* are supplied with a qualifier *massive*; **massive singularities are hidden under an event horizon**- this is a version of CCH that looks plausible.

The formulation in terms of quasi-local quantities is of interest, since it can be pursued further to cover cases in which the standard approach fails. In spherically symmetric geometries (asymptotically flat and in some cosmological models) event horizons must exist if apparent horizons are present [?]. The quasi-local conditions that imply the formation of apparent horizons in spherically symmetric geometries are already known [?], [?]. Using them, one can obtain a number of conditions for the formation of event horizons inside collapsing matter (in asymptotically flat geometries) and in cosmological models.

3. The Cauchy problem.

In the above considerations I have assumed the existence of a global maximal Cauchy surface which possesses a maximal extension at least in the part of a space-

time that is exterior to the apparent horizon and which includes the latter. Let me point out that in standard proofs of the Birkhoff theorem one usually assumes the existence of that part of space-time that is exterior to the collapsing body; that is merely equivalent with my conditions. Nevertheless there exists a possibility to get rid of the assumption. Below I sketch a line of reasoning that should lead to a proof of a version of the global Cauchy problem.

To pursue further we will need the spherically symmetric Einstein equations in electrovacuum. In electrovacuum some of the matter related terms (i. e., j_r, T_μ^μ) of equations (??, ??) vanish. Notice that $K_{ij}K^{ij}$ can be written (due to spherical symmetry) as $\frac{3}{2}(K_r^r)^2$. Below appears the mean curvature p of a sphere as embedded in a hypersurface Σ_t

$$p = \frac{(r\partial_r b + 2b)}{\sqrt{abr}}. \quad (23)$$

The energy density ρ contains only a contribution from the electrostatic field,
 $\rho = \frac{q^2}{8\pi r^4 b^2} = \frac{q^2}{8\pi R^4}.$

The Einstein equations read

$$-\frac{2}{abr^2}\partial_r(r^2\partial_r b) + \frac{(\partial_r b)^2}{2ab^2} + \frac{\partial_r b \partial_r a}{a^2 b} + \frac{2\partial_r a}{a^2 r} - \frac{2\partial_r b}{abr} + \frac{2}{r^2}\left(\frac{1}{b} - \frac{1}{a}\right) = \frac{3}{2}(K_r^r)^2 + 16\pi\rho, \quad (24)$$

$$\partial_r K_r^r = -\frac{3\sqrt{a}}{2}pK_r^r, \quad (25)$$

$$\frac{1}{a^{1/2}br^2}\partial_r(a^{-1/2}br^2\partial_r\alpha) = [\frac{3}{2}(K_r^r)^2 + 8\pi\rho]\alpha, \quad (26)$$

$$\partial_t a = 2\alpha a K_r^r, \quad (27)$$

$$\partial_t K_r^r = \frac{3\alpha}{4}(K_r^r)^2 - \frac{\alpha p^2}{4} - \frac{p\partial_r\alpha}{\sqrt{a}} + \frac{\alpha}{br^2} - 8\pi\alpha\rho. \quad (28)$$

3a. The initial data.

Assume that Σ_0 is a global Cauchy hypersurface. As pointed out at the end of section 1, initial data are determined by the initial distribution of matter and momentum; without loss of generality we can assume that initially $a(r) = b(r)$. If Σ_0 contains an apparent horizon, then a singularity shall develop in the future, therefore the best we can hope to prove is the existence of a solution outside the apparent horizon. We wish to consider the Cauchy problem outside the apparent horizon; this seems to be reasonable, since the apparent horizon moves outward with the speed of light and nothing that happens inside it can casually influence its exterior.

Let Σ_t^{out} be a Cauchy maximal hypersurface that evolves from Σ_0 in the region outside the cylinder enclosed by the apparent horizon. Σ_0^{out} coincides obviously with a corresponding part of Σ_0 , hence the Cauchy data K_r^r and a are fixed. (The function b might be determined from (??) and is not an independent dynamical quantity.) The hamiltonian constraint constitutes an elliptic equation. From asymptotic flatness we have to set $a(\infty) = b(\infty) = 1$, but this condition is not sufficient to ensure the uniqueness of solutions of equation (??) on Σ_t^{out} . However, the asymptotic mass m must be constant on all slices and it is determined by the geometry of Σ_0 . Therefore we must demand that on all Cauchy slices Σ_t^{out} the following asymptotic condition is met

$$\lim_{r \rightarrow \infty} r^2 \partial_r a = \lim_{r \rightarrow \infty} r^2 \partial_r b = -2m \quad (29)$$

.

In what follows we assume that the convergence $\theta' = K_r^r + p$ of ingoing light rays is strictly positive on Σ_0 ; this implies that $\theta' = K_r^r + p$ is strictly positive on all future slices Σ_t^{out} (Appendix 2). If there is an apparent horizon, then $\theta(r) = K_r^r - p$ vanishes at a centered sphere in Σ_t^{out} ; the preceding assumption $\theta' > 0$ implies that $K_r^r > 0$ at the apparent horizon and out of it, since K_r^r does not change sign in electrovacuum (see formula (??)). Now we can conclude that at the outermost apparent horizon and out of it we must have $p > 0$, i.e., there is no minimal surface in Σ_t^{out} . Let $r = r_t$

be a position of the outermost apparent horizon. We may invoke the calculation performed in the previous section, which led to the equation (??). (The calculation bases on the assumption that $a = b = \phi^4$ but it is only a technical condition and there is no any loss of generality.) At the apparent horizon (??) yields

$$m - [\sqrt{\frac{S}{16\pi}} + q^2 \sqrt{\frac{\pi}{S}}] = 0. \quad (30)$$

In electrovacuum there are two solutions of the equation (??),

$$r_t \phi^2(r_t)|_2^1 = m_-^\pm \sqrt{m^2 - q^2}; \quad (31)$$

taking into account the above conditions we have to choose

$$\phi(r_t) = \sqrt{\frac{1}{r_t}(m + \sqrt{m^2 - q^2})} \quad (32)$$

since otherwise there could exist a minimal surface at some $r > r_t$. The hamiltonian constraint is now supplied with the standard Dirichlet boundary conditions $\phi(r_t), \phi(\infty) = 1$ on Σ_t^{out} and it is easy to prove that there is a unique solution.

Thus, under the above conditions fixing the asymptotic mass m uniquely determines the conformal factor ϕ (and, consequently, a, b , if a relation between the two functions is determined on an initial slice) at the surface of the apparent horizon. But m is determined by the initial geometry Σ_0 , hence we have no any freedom left in specifying the solutions of the hamiltonian constraint in the exterior region.

The corresponding boundary problem for the lapse equation (??) contains, however, an arbitrariness. The condition

$$\alpha(\infty) = 1 \quad (33)$$

does not specify uniquely a solution of (??); we still can impose

$$\partial_r \alpha = f(t) \quad (34)$$

at the surface of an outermost apparent horizon. I assume that the function $f(t)$ is smooth and strictly positive. We have defined the exterior Cauchy problem (i. e.,

with data on Σ_t^{out}) as a restriction of the global Cauchy problem with data on Σ_t . There is an obvious loss of information during such a restriction, since we do not control the collapse of matter fields that are enclosed inside the apparent horizon. The arbitrariness in choosing $f(t)$ corresponds to our unawareness about the full state of the collapsing system.

In summary, the initial value problem of the electrovacuum Einstein equations outside the apparent horizon can be determined by prescribing the asymptotic mass m , a relation $a(r, t = 0) = b(r, t = 0)$, an initial datum K_r^r at a time $t = 0$ and a condition (??) for the lapse function.

3b. The exterior Cauchy solution exists globally.

Theorem 2. Let $H_s(\Sigma_t^{out})$ be a Sobolev space of functions defined on Σ_t^{out} . Let $\Sigma_{t=0}$ be a global maximal Cauchy surface with initial data $a(t = 0) = b(t = 0)$ such that $\partial_r a(r, t = 0) \in H_2(\Sigma_{t=0})$, $K_r^r(t = 0) \in H_2(\Sigma_{t=0})$ generated by a given initial distribution of matter. Assume that the convergence $\theta'(S)$ of the ingoing light rays is strictly positive for any centered sphere S on Σ_0^{out} and that there exists an apparent horizon that is placed in vacuum or in electrovacuum. Let the lapse function satisfy the boundary conditions (??, ??) with $f(t) \geq 0$ at the outermost apparent horizon. Then there exists a unique solution of the global Cauchy problem in the region exterior to the apparent horizon, including the apparent horizon itself.

Sketch of the proof.

There exist theorems [?] from which one infers the existence of a solution $(K_r^r \in H_s(\Sigma_t), \partial_r a \in H_s(\Sigma_t) \text{ or } K_r^r, \partial_r a \in H_s^{loc}(\Sigma_t))$ for $s \geq 2$ and sufficiently short intervals of time. The lower bound on the index s is due to the Schauder ring property [?] which is fulfilled in three spatial dimensions if $s \geq 2$. Assuming the local existence, I will estimate (Step 1) the pointwise growth of $a(t)$, $K_r^r(t)$, $\partial_r K_r^r$, p , $\partial_r p$, $\partial_r \alpha$. Using those estimates and the method of energy estimates [?] one can show (Step

2) that Sobolev norms in question $(K_r^r \epsilon H_s(\Sigma_t^{out}), \partial_r a \epsilon H_s(\Sigma_t^{out}))$ do not blow up in a finite time t , thus accomplishing the final goal.

Step 1. The L_∞ estimates.

In electrovacuum the right hand side of (??) is nonnegative. Invoking the maximum principle and using the boundary condition that the lapse satisfies at the apparent horizon we conclude that

$$\partial_r \alpha \geq f(t) \geq 0 \quad (35)$$

everywhere in Σ_t^{out} . As pointed out in the subsection (3a), if $\theta' > 0$ then

$$p > K_r^r > 0 \quad (36)$$

everywhere in Σ_t^{out} (but at spatial infinity both functions vanish, $K_r^r = O(1/r^3)$ and $p = 0(1/r)$). Notice also, that

$$a(r, t) = a(r, 0) + 2 \int_0^t \alpha K_r^r ds \geq a(r, 0) = b(r, 0) \geq b(r, 0) - 2 \int_0^t \alpha K_r^r ds = b(r, t) \quad (37)$$

.

Using the inequalities (??, ??) one gets from (??) and (??)

$$(\partial_t + \frac{\alpha}{\sqrt{a}} \partial_r) K_r^r \leq \frac{1}{br^2} \leq \frac{1}{R_{AH}^2}, \quad (38)$$

where I used $\alpha \leq 1$ and the obvious fact (since $p > 0$ on Σ_t^{out}) that the areal radius $R = \sqrt{br}$ is bounded from below by the areal radius of the apparent horizon $R_{AH} = m + \sqrt{m^2 - q^2}$. Let S^* be an intersection of Σ_0^{out} with a light cone passing through a sphere of a coordinate radius r in Σ_t^{out} and let r^* be a coordinate radius of S^* .

$K_r^r(t, r)$ is bounded from above,

$$K_r^r(r, t) \leq K_r^r(r^*, 0) + \frac{t}{R_{AH}^2}. \quad (39)$$

The equation (??) can be solved on each slice Σ_t^{out} to give

$$K_r^r(r, t) = \frac{C(t)}{R^3}. \quad (40)$$

At the apparent horizon $p = K_r^r$ and $\theta' = p - K_r^r$ is positive during the collapse (Appendix 2) if it is positive on Σ_0^{out} ; this implies that K_r^r is positive at the horizon and from (??) also $C(t)$ is positive.

The extrinsic curvature K_r^r decreases on a fixed slice Σ_t^{out} and achieves its largest value at the apparent horizon. The chain of inequalities

$$\frac{C(t)}{R^3} \leq \frac{C(t)}{R_{AH}^3} \leq \frac{C(0) + tR_{AH}}{R_{AH}^3} \quad (41)$$

leads to the final estimations

$$C(t) \leq C(0) + tR_{AH}, \quad K_r^r(r, t) \leq \frac{C(0) + tR_{AH}}{R^3}. \quad (42)$$

(??) and (??) imply an estimation

$$\partial_t a(r, t) = 2a(r, t)\alpha(r, t)K_r^r(r, t) \leq a(r, t)2\frac{C(0) + tR_{AH}}{R_{AH}^3} \quad (43)$$

which in turn gives the following estimation of the metric coefficient $a(r, t)$:

$$a(r, t) \leq a(r, 0)e^{\frac{2tC(0) + t^2 R_{AH}}{R_{AH}^3}}. \quad (44)$$

A straightforward calculation gives

$$\partial_t p(r, t) = -\frac{\partial_r \alpha(r, t)}{\sqrt{a}} K_r^r(r, t) + \frac{\alpha K_r^r}{2} p \leq \frac{\alpha K_r^r}{2} p \leq \frac{C(0) + tR_{AH}}{2R_{AH}^3} p; \quad (45)$$

above I employed the inequalities $\partial_r \alpha \geq 0$, $\alpha \leq 1$ and (??).

One can show also that

$$\partial_r p(r, t) = -\sqrt{a}(8\pi\rho + \frac{3}{4}(K_r^r(r, t))^2 + \frac{3}{4}p^2 - \frac{1}{br^2}). \quad (46)$$

(??), (??) and the previous estimates yield

$$(\partial_t + \frac{\alpha}{\sqrt{a}}\partial_r)p = 8\pi\alpha(\frac{j_r}{\sqrt{a}} - \rho) - \frac{\partial_r \alpha}{\sqrt{a}} K_r^r - \frac{\alpha}{4}[\theta^2 + 2(K_r^r)^2 + 2p^2] + \frac{\alpha}{R^2} \leq \frac{1}{R^2}, \quad (47)$$

which gives finally

$$p(r, t) \leq \sup[p(r, t = 0)] + \frac{t}{R_{AH}^2}. \quad (48)$$

A combination of (??, ??) and (??) together with the momentum constraint (??) allows one to obtain a bound for $\partial_r K_r^r(r, t)$:

$$\left| \frac{\partial_r K_r^r(r, t)}{\sqrt{a(r, t)}} \right| \leq \frac{3(C(0) + tR_{AH})}{R^3} (\sup[p(r, 0)] + \frac{t}{2R_{AH}^2}). \quad (49)$$

The energy density is positive and bounded from above by $\frac{q^2}{8\pi R_{AH}^4}$; using the preceding information one easily infers that $|\partial_r p|$ must be uniformly bounded on all slices Σ_t^{out} by $C_1 t^2 + C_2 t + C_3$ where C_1, C_2, C_3 are constants depending only on the initial data.

In order to get estimations of $\partial_r \alpha(r, t), \partial_r^2 \alpha(r, t)$ one should analyze the lapse equation (??) using the above information about the evolution of the extrinsic curvature. A lengthy and not particularly illuminating calculation gives a bound

$$\left(\frac{\partial_r \alpha}{\sqrt{a}} \right)(r, t) \leq f(t) + \frac{4\pi q^2}{RC(t)} + \frac{3\pi C(t)}{R^2 R_{AH}}. \quad (50)$$

The right hand side of (??) is finite since $C(t)$ is strictly greater than 0 in any finite time t (Appendix 2) and satisfies (??).

In summary, the following estimates hold true

$$\|X\|_{L_\infty(\Sigma_t^{out})} \leq F(t) \|X\|_{L_\infty(\Sigma_0^{out})}, \quad (51)$$

where X is any of the functions $(a, K_r^r, \partial_r K_r^r, p,)$ and $F(t)$ is a positive function that remains bounded for arbitrarily large but finite values of its argument. The lapse α does not exceed 1, $\partial_r \alpha$ must satisfy (??) and $|\partial_r p|$ is bounded by a quadratic function of $\|X\|_{L_\infty(\Sigma_t^{out})}$.

Step 2. The integral estimates.

In order to prove the existence of a global solution one has to show that the Sobolev norm $\|K_r^r\|_{H_2(\Sigma_t^{out})} + \|\partial_r a\|_{H_2(\Sigma_t^{out})}$ remains bounded for any finite time t .

This could be done explicitly, by pursuing the above calculation in order to get pointwise estimates for all quantities in question and then proving the required integrability. I will choose a way that is probably less economic in this particular case but offers a chance for generalization.

Let $\mathbf{X}$ be a vector having 6 components $\mathbf{X}_i = \partial_r^i a, \mathbf{X}_{i+3} = \partial_r^{i-1} K_r^r, i = 1, 2, 3$. Define

$$H(t) = (\|K_r^r\|_{H_2(\Sigma_t^{out})}^2 + \|\partial_r a\|_{H_2(\Sigma_t^{out})}^2). \quad (52)$$

A simple but laborious calculation shows that

$$\frac{d}{dt}H = \sum_{i,j=1}^6 \int_{\Sigma_t^{out}} dV \mathbf{X}_i \mathbf{X}_j f_{ij} + \sum_{i=1}^6 f_i X_i - 4\pi\alpha R^2 \sum_{i=1}^6 X_i^2 |_{AH}, \quad (53)$$

where f_i, f_{ij} are certain polynomials of finite order that depend on $\sqrt{a}, b, K_r^r, \partial_r K_r^r, p, \partial_r p, \partial_r \alpha, \alpha, \rho$. The functions f_i are square integrable. The crucial point is that the right hand side of (53) is bilinear in $\mathbf{X}_i$. Direct differentiation of $\|K_r^r\|_{H_2(\Sigma_t^{out})}^2 + \|\partial_r a\|_{H_2(\Sigma_t^{out})}^2$ with respect t and the use of evolution equations (54, 55) gives also some trilinear terms but manipulating with equations (54- 55) and (56) finally yield the above equation.

The right hand side of (53) is bounded from above by

$$\sup\{\|f_{ij}\|_{L_\infty(\Sigma_t^{out})}\} \sum_{i=1}^6 \int_{\Sigma_t^{out}} dV \mathbf{X}_i \mathbf{X}_i + \sum_{i=1}^6 [\int_{\Sigma_t^{out}} dV f_i^2]^{1/2} \sum_{i=1}^6 [\int_{\Sigma_t^{out}} dV \mathbf{X}_i \mathbf{X}_i]^{1/2}. \quad (54)$$

Taking into account estimates of the preceding subsection, one obtains the inequality

$$\partial_t H \leq C_2(t)H + C_3(t), \quad (55)$$

where C_2, C_3 are exponentially bounded functions of coordinate time t with coefficients depending only on initial data. (55) readily implies

$$H(t) \leq e^{\int_0^t ds C_2(s)} \int_0^t ds C_3(s) e^{-\int_0^t ds C_2(s)} + H(t=0) e^{\int_0^t ds C_2(s)}. \quad (56)$$

thus proving the existence of a solution for any finite time t . That ends the proof of Theorem 2.

Remarks. The Cauchy problem should be investigated in *weighted Sobolev spaces* [?] instead of Sobolev spaces. The hypersurfaces Σ_t^{out} are noncompact and one should incorporate suitable falloff conditions at spatial infinity; weighted Sobolev spaces include them automatically, in contrast with the standard Sobolev spaces. It is easy, however, to adapt the above proof to work with $H_{s,\delta}$ instead of H_s and I will not discuss this point.

4. Final comments.

The present investigation bases on the (1+3) splitting of spacetime that is smooth (initially and possibly also globally, modulo a region of compact support). The world time t can be used globally to parametrize casually related occurrences. The proof that in electrovacuum event horizons coincide with apparent horizons is done with only minimal reference to specific properties of spherically symmetric geometries. Obviously, the existence of the maximal slicing requires a proof [?], but the presence or absence of spherical symmetry is probably of no great significance for the validity of maximal slicings. The place where the assumption of spherical symmetry plays important role is the proof of the identity (??) but it is quite likely that (??) survives (in the form of the Penrose-Gibbons inequality) also in nonspherical geometries. Alternatively, one can use the Raychaudhuri equation (??) in order to prove the local confining property of apparent horizons, which should be of help in proving the existence of the global Cauchy solution.

The global Cauchy problem, however, poses a serious obstacle in making a significant progress in proving the Cosmic Censorship Hypothesis.

The application of the above ideas to a more general class of spherically symmetric

geometries of collapsing systems will be reported elsewhere [?].

Acknowledgements. I wish to thank Niall O'Murchadha for many discussions and valuable remarks, that greatly contributed to my understanding of problems discussed here. This work was supported in part by the Polish Government Grant KBN 2526/93.

Appendix 1

Static Einstein equations reduce to three equations (??), (??) and (??), of which only two are independent. Equation (??) gives the spatial metric, which in a gauge $a(r) = b(r) = \phi^4$ coincides with the solution (??). Inserting this solution into (??) gives (notice that $K_r^r = 0$ and $c = \frac{m^2 - q^2}{4}$)

$$\begin{aligned} \partial_r \ln \alpha &= \frac{\sqrt{a}}{pbr^2} - \frac{p\sqrt{a}}{4} - \frac{q^2\sqrt{a}}{b^2r^4p} = \\ (\text{here } p &= \frac{2(2\partial_r\phi r + \phi)}{\phi^3r}) \\ &= \frac{mr^2 + 4cr + mc}{(r^2 - c)(r^2 + mr + c)}. \end{aligned}$$

The last equation is solved by (??).

Appendix 2

Lemma. Under conditions stated in Theorem 2, if $\theta' = p + K_r^r$ is positive on Σ_0 then it must be positive in the all future slices Σ_t^{out} .

Proof. Assume the contrary, i. e., that there exists a Cauchy (external) hypersurface Σ_t^{out} such that somewhere on it θ' crosses through zero.

The evolution of θ' is given by the equation

$$\begin{aligned} (\partial_t - \frac{\alpha}{\sqrt{a}}\partial_r)\theta' &= \\ (-\partial_r\alpha/\sqrt{a} + \alpha K_r^r)\theta' + \alpha\theta'^2/2. \end{aligned}$$

From this equation one infers that the surface with vanishing convergence θ' moves inward with the velocity of light when immersed in vacuum (Section 2). Therefore it must exist in all preceding Cauchy slices and in particular in the initial hypersurface Σ_0^{out} . This gives a contradiction which proves our claim.

Corollary. Assume that Σ_0 contains an apparent horizon. Under conditions of the preceding lemma, $C(t)$ (and, consequently, K_r^r) must be strictly positive for any finite time t .

Proof. From the results of Section 2, the apparent horizon propagates to the future. If $C(t)$ was equal to zero on a slice Σ_t^{out} for some time t , then both θ and θ' would vanish at an apparent horizon, which would imply (due to the above lemma) the existence of a white hole in Σ_0 , contrary to the assumption that $\theta' > 0$ in Σ_0 .

REFERENCES

- [1] R. Penrose, Riv. Nuovo Cimento **1**, 252 (1969).
- [2] The only global result is that of D. Christodoulou, S. Klainerman, Acta Mathematica, (1991) who prove the existence of space-times that are almost Minkowskian.
- [3] D. Christodoulou, Commun. on Pure Appl. Math. **44**, 399(1991).
- [4] R. Penrose, Phys. Rev. Lett. **14**, 252 (1969).
- [5] W. Israel, Phys. Rev. Lett. **56**, 86(1986).
- [6] P. Bizoń, E. Malec and N. O’ Murchadha, Phys. Rev. Lett. **61**, 1147(1988); Class. Quantum Gravity **7**, 1953(1990).
- [7] U. Brauer and E. Malec, Phys. Rev. D **45**, R1836(1992); E. Malec and N. O’ Murchadha, Phys. Rev. D **47**, 1454(1993); T. Zannias, Phys. Rev. D **45**, 2998(1992), reformulated results of [?] using nonmaximal slicings; S. Hayward, Class. Quantum Grav. **9**, L135(1992) used 2+2 slicings of space-time.
- [8] e. g., C. W. Misner, J. A. Wheeler and K. S. Thorne, *Gravitation* Freeman 1973.
- [9] S. Hawking and G. F. R. Ellis, The large scale structure of space-time, Cambridge University Press 1973.
- [10] I am convinced that this form of the electrovacuum (Reissner-Nordstroem) solution of Einstein-Maxwell equations has been obtained hitherto – its space part is well known, see [?] – but I was not able to find any reference in which the lapse function is given.
- [11] G. Gibbons, p. 194 in Global Riemannian Geometry, ed. N. J. Willmore and N. J. Hitchin, Elli Horwood, New York 1984.
- [12] R. Penrose, p. 631 in Seminar on Differential Geometry Princeton University Press 1982.
- [13] J. Isenberg, N. O’ Murchadha and J. W. York Phys. Rev. D **13**, 1532(1976).